

2-22-2012

# Effects of road salt on photosynthetic and enzyme activity of stream biofilms

Leah Cook

Follow this and additional works at: <http://commons.emich.edu/theses>



Part of the [Biology Commons](#)

---

## Recommended Citation

Cook, Leah, "Effects of road salt on photosynthetic and enzyme activity of stream biofilms" (2012). *Master's Theses and Doctoral Dissertations*. 410.

<http://commons.emich.edu/theses/410>

This Open Access Thesis is brought to you for free and open access by the Master's Theses, and Doctoral Dissertations, and Graduate Capstone Projects at DigitalCommons@EMU. It has been accepted for inclusion in Master's Theses and Doctoral Dissertations by an authorized administrator of DigitalCommons@EMU. For more information, please contact [lib-ir@emich.edu](mailto:lib-ir@emich.edu).

EFFECTS OF ROAD SALT ON PHOTOSYNTHETIC AND ENZYME ACTIVITY OF  
STREAM BIOFILMS

BY

Leah J. Cook

Thesis

Submitted to the Department of Biology

Eastern Michigan University

In partial fulfillment of the requirements

for the degree of

MASTER OF SCIENCE

in

Ecology and Organismal Biology

Thesis Committee:

Steven N. Francoeur, PhD, Chair

Peggy Liggitt, PhD

Gary Hannan, PhD

February 22, 2012

Ypsilanti, Michigan

THESIS APPROVAL FORM

EFFECTS OF ROAD SALT ON PHOTOSYNTHETIC AND ENZYME ACTIVITY OF  
STREAM BIOFILMS

BY

Leah J. Cook

APPROVED:

\_\_\_\_\_  
Steven N. Francoeur, PhD  
Thesis Chair

\_\_\_\_\_  
Date

\_\_\_\_\_  
Peggy Liggitt, PhD  
Committee Member

\_\_\_\_\_  
Date

\_\_\_\_\_  
Gary Hannan, PhD  
Committee Member

\_\_\_\_\_  
Date

\_\_\_\_\_  
Marianne Laporte, PhD  
Department Head

\_\_\_\_\_  
Date

\_\_\_\_\_  
Deborah de Laski-Smith, PhD  
Dean of the Graduate School

\_\_\_\_\_  
Date

## **DEDICATION**

To the three pillars in my life: God, my husband, and my children.

“For I know the plans I have for you,” declares the LORD,

“plans to prosper you and not to harm you, plans to

give you hope and a future” Jeremiah 29:11.

DeWayne, you are an amazing, supportive husband

who continues to push me to do my best in everything.

I love you and thank you so much for always being there.

Charlotte and Lukas, you are the legacy of all my

hard work. “Mommy’s homework” is done.

## ACKNOWLEDGEMENTS

I am forever grateful for the patience and direction provided by Dr. Steven N. Francoeur in completing my Master's research. I am thankful for his friendship and his guidance during this process. Similar gratitude and appreciation are owed to Dr. Peggy Liggitt and Dr. Gary Hannan for their encouragement, editorial comments, and critical advice throughout this process. I wish to thank, Kevin Johnson, Mark Schaecher, Audrey Johnson, and Sarah Whorley, who helped with the very cold field aspect of this project. I grateful to Audrey Johnson and Sarah Whorley for all of their expertise in analyzing nutrient data. Fungal data would not have been possible without the efforts of Dr. Kevin A. Kuehn. Without all of the aforementioned people, my thesis research would not have been completed.

I would also like to thank the Washtenaw County Office of the Water Resources Commissioner for allowing access to Paint Creek where this study was conducted, and the Huron River Watershed Council for providing local stream conductivity data. This research was supported, in part, by funding from the Center for Aquatic Microbial Ecology and the Graduate School of Eastern Michigan University. Additional support was provided by scholarship awards from the Meta Hellwig Scholarship Fund of the Department of Biology of Eastern Michigan University.

## **ABSTRACT**

Road salt application and salt-laden run-off are common, yet little information exists about its influence on periphyton. I measured extracellular enzyme activity and photosynthetic activity of natural winter stream periphyton communities under various short-term laboratory salt exposures. The results of this study suggest that short-term exposure to increased concentrations of salt did not affect extracellular enzyme activity but could temporarily reduce the photosynthetic activity of the periphyton. Further work will be necessary to understand the long-term effects of increased salt concentrations on periphyton and extracellular enzyme activity.

## TABLE OF CONTENTS

APPROVAL .....	ii
DEDICATION .....	iii
ACKNOWLEDGEMENT .....	iv
ABSTRACT.....	v
LIST OF TABLES .....	vii
LIST OF FIGURES.....	viii
INTRODUCTION.....	1
MATERIALS AND METHODS.....	7
RESULTS.....	13
DISCUSSION.....	16
REFERENCES.....	22

## LIST OF TABLES

<u>Table</u>		<u>Page</u>
1	Environmental conditions at the Paint Creek field site. Values are means ( $\pm$ 1 SD) of six measurements made from 4 January – 13 March 2007). Asterisk indicates water samples were taken from stream and water temperature and conductance were measured upon returning to lab. n.d. = not determined .....	26
2	Mean ( $\pm$ 1 SE) algal biomass, fungal biomass, and bacterial abundance of periphyton communities used for the three experiments. n.d. = not determined.....	27

## LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
1	Conductivity measurements ( $\mu\text{S cm}^{-1}$ , $n = 1456$ , 1995-2006) from Huron River tributaries in Washtenaw County, MI and surrounding region. Data are sorted highest to lowest, and are courtesy of the Huron River Watershed Council (HRWC.org).....	28
2	Effects of conductivity on dark-adapted photosystem II yield (Fv/Fm) in experiment one ( $p < 0.001$ ). Values are means $\pm 1$ SE ( $n = 7$ ).....	29
3	Effects of conductivity on dark-adapted photosystem II yield (Fv/Fm) in experiment three. Photosystem II yield measurements means were significantly different before a rinse ( $p < 0.001$ ), but not significantly different following the rinse at 6 hours and before 6.5 hours ( $p = 0.061$ ).....	30
4	Mean extracellular enzyme activity ( $\text{nmol cm}^{-2} \text{h}^{-1}$ at ambient stream water ( $2009.2 \mu\text{S cm}^{-1}$ ) and salt-amended stream water ( $\sim 60,000 \mu\text{S cm}^{-1}$ )) .....	31

## INTRODUCTION

Winter roads are often maintained by the application of some form of chemical deicer, often road salt. Removing snow and salting of roads during wintry weather is used to increase the safety of commuters. Salt lowers the freezing point of water, thus preventing ice formation and encouraging melting (Oškinis & Kasperovičius, 2005). Maintenance of winter roads by applying salt was first introduced in the 1930s and has been used more frequently since the 1960s (Paschka et al., 1999). Several salts are used on roads. The most common are calcium chloride and sodium chloride (Environment Canada, 2001; Ramakrishna & Viraraghavan, 2005). Additives may be used to increase the effectiveness of the salt (Novotny et al., 2008). In Washtenaw County, Michigan, the typical annual use of salt by the Washtenaw County Road Commission is 10,000 to 25,000 tons of salt applied on its ~ 2,000 miles of roadways (<http://www.wcroads.org/services/operations/saltusage.htm>, last accessed 04-30-2010). For example, in the 2006/2007 winter season, the Washtenaw County Road Commission used 16,837 tons of salt. These figures do not include road salt applications by the numerous municipalities within Washtenaw County, nor do they include applications by private individuals on sidewalks, driveways, and parking lots.

The clearing of snow from roadways, sidewalks, and driveways contributes to the salt entering nearby habitats (Environment Canada, 2001). Furthermore, aquatic habitats near roadways or other impervious surfaces acquire runoff containing a greater concentration of deicing chemicals (Novotny et al., 2008; Paschka et al., 1999; Environment Canada, 2001). High salt concentrations enter into streams and lakes through snowmelt runoff (Oškinis & Kasperovičius, 2005). This runoff can be either

direct (surface flow into a nearby surface water) or indirect (runoff that is detained or infiltrated into the aquifer then into nearby water systems). Increased conductivity in streams during winter months resulting from salt applications is common (Oškinis & Kasperovičius, 2005). Furthermore, some aquatic habitats do not have the ability to dilute the salts, which can influence the water chemistry of the system (Environment Canada, 2001). Lakes can experience a seasonal cycle of chloride related to road salt applications (Novotny et al., 2008), and salt runoff can alter water densities in lakes, leading to salt-induced lake stratification (Ramakrishna & Viraraghavan, 2005; Dodds & Whiles, 2010). Lakes in Washtenaw County have experienced similar impacts (Judd et al., 2005), and winter stream conductivities can be greatly elevated (Fig. 1).

Adding salt to an environment can influence the environment and its aquatic organisms. Increased salt within the environment can change the density of water and reduce the solubility of other molecules. Increased water density influences lake stratification (Judd et al., 2005; Novotny et al., 2008). In addition to impacting aquatic habitats, increased salt in the environment can influence ions found in the soil (Environment Canada, 2001).

Organisms that live in or near aquatic systems vary in their tolerances to salt. A wide variety of organisms have been studied in relation to their exposure to salt in the field and in the laboratory. These organisms include fish, zooplankton, and benthic invertebrates where laboratory studies investigated chloride toxicity in short term exposures (Environment Canada, 2001). Further studies using amphibians and road salt suggest that aquatic organisms react negatively to increased salt concentrations. Sanzo and Hecnar (2006) observed the influence of sodium chloride on larval wood frogs (*Rana*

*sylvatica*). They found that salt concentrations lowered survivorship and reproductive fitness. The frogs also displayed more physical abnormalities with increased concentrations of salt. In addition, Denoël et al. (2010) concluded that tadpoles (*Rana temporaria*) exposed to high salt concentrations had a slower movement speed than control tadpoles.

Macroinvertebrates also have been used to study the effects of salts on aquatic systems. Researchers have classified macroinvertebrates as tolerant or non-tolerant to salt (Williams et al., 1999). Blasius and Merritt (2002) found that macroinvertebrate community composition and mortality were affected by short-term exposure to elevated salt concentrations; some macroinvertebrates were more tolerant than others of higher concentrations of salt.

Diatom assemblages can indicate changes in conductivity of aquatic habitats. Indices have been created to reflect the conductivity ranges of ecosystems using diatoms (Blinn, 1993; Clavero et al., 2000). Increasing conductivity in established mesocosms decreased species richness and diversity in diatom assemblages. Decreased diatom diversity and species richness in relation to increased conductivity were also observed by Blinn (1993), who collected diatoms from 63 North American lakes. In addition, the researcher found a significant relationship between particular anions and diatom species in low and high conductivity environments. The presence of salt may hinder the growth of organisms or change the community composition within ecosystems (Herbst & Blinn, 1998). Growth of diatoms is generally reduced in the presence of higher salinity (Saro & Fritz, 2000). When excess salinity is introduced to an environment, organisms often need to allocate energy for self-maintenance rather than growth (Calow, 1991). Photosystem II

in algae can be inhibited by the presence of excess salt (Allakhverdiev et al. 2000; Lu & Zhang, 2000; Sudhir et al., 2005).

In addition, algae exposed to salt can exhibit decreased photosystem II activity and decreased oxygen production (Allakhverdiev et al., 2000; Lu & Zhang, 2000; Sudhir et al., 2005). The cyanobacterium *Synechococcus* sp. provides an example of how a photosynthetic aquatic organism adapts to ionic and osmotic stressors. Ionic and osmotic stressors have been studied to determine their influence upon Photosystem I (PSI) and Photosystem II (PSII) in this organism (Allakhverdiev et al., 2000). Cyanobacteria were exposed to 0.5M NaCl, and oxygen production due to PSII was measured. A rapid decrease of oxygen production was observed after 0.5, 1, or 5 hours exposure to 0.5M NaCl. Following the shortest exposure, washing cells and returning them to a normal culture medium resulted in complete recovery of photosynthetic activity. After the 1-hour NaCl incubation, the cells recovered partially when returned to normal culture medium. After the 5-hour NaCl incubation, the cells showed no recovery when washed and returned to normal culture medium. Similar experiments using exposure of cells to non-ionic sorbitol indicated that both ionic and osmotic effects can contribute to reduced oxygen production caused by exposure to salts such as NaCl (Allakhverdiev et al., 2000).

Changing salinity is a form of osmotic stress that changes the osmotic equilibrium (Dodds & Whiles, 2010). Organisms exposed to higher salinity levels also experience osmotic water loss. If equilibrium cannot be established, the pressure from the osmotic gradient will dehydrate the cells and they will collapse (Dodds & Whiles, 2010). In addition, the increased ion concentrations can influence electron transport activities and thylakoid membrane protein function. The inhibition of protein function suggests the

potential for inhibition of PSII activity (Sudhir et al., 2005). Furthermore, an increase in salinity may cause metals to be more toxic by influencing rates of metal adsorption and uptake (Borchardt, 1996)

The rate and duration of salt inputs to aquatic environments may be moderated by soils. Negatively charged chloride ions rapidly pass through the negatively-charged soil (Environment Canada, 2001). This will contribute to increased salinity in the soil. The positive sodium ions will also move through the soil, but at a slower rate, and can displace calcium and magnesium cations. Long-term salt application may result in soils with elevated salinity; such soils will leach salt into nearby pools or streams, causing elevated aquatic salt concentrations at various times of the year (Ramakrishna & Viraraghavan, 2005; Kelly et al., 2008).

Periphyton is the community of microorganisms (e.g., algae, bacteria) that grows attached to surfaces in aquatic environments. Benthic algae are an important component of periphyton in streams, contributing to the base of the food web and serving as water quality indicators (Hauer & Lamberti, 1996). Benthic algae are important primary producers in streams and serve as a food source for stream invertebrates (Borchardt, 1996). Furthermore, they are necessary as chemical modulators, stabilizers of substrata, and providers of habitats for other organisms (Borchardt, 1996). There are several factors that could limit benthic algae within a stream including “light, temperature, current, substrate, scouring by floods, water chemistry and grazing” (Allan & Castillo, 2007).

Periphyton help transfer organic matter through the aquatic habitat by utilizing extracellular enzymes to hydrolyze organic molecules into smaller ones for microbial use (Pohlen et al., 2010; Allan & Castillo, 2007). Both bacterial and algal components of

periphyton communities produce and secrete extracellular enzymes. Enzyme activity can indicate a shift in physiological state of algae and bacteria. Ratios of the activities of N-, P- and C-acquiring extracellular enzymes can be used to infer whether microorganisms are N-, P-, or C-limited (Rier et al., 2011; Sinsabaugh et al., 2010). The release of such enzymes can also influence dissolved organic carbon concentrations within a stream (Allan & Castillo, 2007), and enzyme activity is related to decomposition rates of detritus (Sinsabaugh et al., 1994; Sinsabaugh & Linkins, 1993).

Little is known about the influence of salinity arising from road salt application on the functioning of stream periphyton. Studies have shown that high salinities have an effect on the overall functioning of stream biota within a stream (Silva et al., 2000). Photosynthesis of freshwater algae can be enhanced by moderate ion concentrations ( $< \sim 5800 \mu\text{S cm}^{-1}$ ) and reduced by higher levels ( $\sim 10,200 \mu\text{S cm}^{-1}$ ; Silva & Davies, 1999). Enzyme activity could be affected by chemical conditions within the matrix of the biofilm (Rier et al., 2007). However, very little research has focused on the influence of conductivity on enzyme activity. In natural systems, elevation of ionic strength via addition of humic acids inhibited extracellular enzyme activity (Dudley & Churchill, 1995).

The purpose of this study is to determine if periphyton photosynthetic activity and extracellular enzyme activity could be influenced by road salt run-off. Thus, the hypothesis was that a reduction of periphytic photosynthetic and enzymatic activity would occur with increased conductivity. This was tested by conducting laboratory experiments that exposed natural winter periphyton communities to realistic salt concentrations.

## MATERIALS AND METHODS

### *Study Site*

The study was conducted in Paint Creek (N42.21502, W83.63494), just downstream of the Paint Creek wetland retention basin within the Stony Creek (Michigan, USA) watershed. Water temperature, specific conductance (YSI model 63), depth, and photosynthetically-active radiation (LI-Cor model 189) above the water surface were recorded during sampling visits. Stream water was also collected for experimental use and measurements of total phosphorus (TP), ammonia ( $\text{NH}_4^+$ ), nitrate and nitrite ( $\text{NO}_3^- + \text{NO}_2^-$ ) were made. Water samples for nutrient assays were frozen until analysis (method of Lind, 1985) and samples for  $\text{NH}_4^+$ , and  $\text{NO}_3^- + \text{NO}_2^-$  analyses were filtered (0.7- $\mu\text{m}$  pore size glass-fiber filters) prior to freezing.

### *Experiment One (February 6, 2007)*

Glass fiber filters were prepared by sandwiching the filters between two plexiglass plates, one of which contained openings through which periphyton colonized the filters (see Kahn & Wetzel, 1999; Francoeur & Wetzel, 2003). The plexiglass plates were wired to bricks and placed on the streambed. After four weeks of periphyton growth (February 6, 2007), the glass fiber filters were retrieved and returned to the lab. Care was taken to avoid losses of periphyton during collection and transport.

In the lab, four sections were cut from the filters using a cork borer (#10, 1.5 cm diameter). These samples were preserved in 5% glutaraldehyde and stored at 4°C in darkness for later microscopy to determine algal community composition. Preliminary scans indicated that all algae present were diatoms, so samples were cleaned and mounted

for microscopy (method of Carr et al., 1986). A minimum of 100 diatoms per sample were identified and enumerated (1000×).

Three replicate sections (each ~176.7mm<sup>2</sup>) of glass fiber filters were collected and frozen for subsequent chlorophyll analysis by method of Biggs and Kilroy (2000). Chlorophyll was extracted in 90% ethanol (80°C for 5 minutes), followed by overnight (4 °C, darkness) steeping. Samples were then lightly centrifuged to ensure settling of particulate matter, and chlorophyll content was quantified from spectrophotometric absorbance at 665nm and 750nm, with acidification to correct for pheopigments.

The collected stream water was used to generate two salt treatments: ambient (1637.2 μS cm<sup>-1</sup>, unamended site water) and 35,320 μS cm<sup>-1</sup> (site water amended with reagent-grade NaCl). Seven replicate sections (~176.7 mm<sup>2</sup>) of glass fiber filter substrata were subjected to each salt treatment. The substrata were immersed in 5.0 mL of the treatment solution for 30 minutes before the first measurement (0 hours) of dark-adapted photosystem II yield (Walz Diving PAM, see Consalvey et al., 2005). Additional dark-adapted yield measurements were collected at 3 hours, 6 hours, and 24 hours. For the entire course of the experiment, all samples were held in darkness at 5 °C.

#### *Experiment Two (March 6, 2007)*

Naturally-senesced *Typha* litter was collected from the Paint Creek wetland < 100 m upstream of the sampling site and placed in floating mesh trays (see Francoeur et al., 2006). After eight weeks of periphyton growth (March 6, 2007), the *Typha* trays were retrieved and returned to the lab. Care was taken to avoid losses of periphyton during transport.

Algal biomass (as chlorophyll a, n = 3) and community composition (n = 4) were quantified as in Experiment 1, with the exception that known lengths of *Typha* litter were cut for quantitative analyses (see Francoeur et al., 2006). An additional four litter sections were frozen for subsequent analysis of fungal biomass (method of Francoeur et al., 2006; Table 2). Dry mass of the litter and associated microbial communities was measured by drying litter sections (105 °C for 24 h) in pre-weighed, pre-ashed aluminum pans, cooling in a dessicator, and weighing. Ash-free dry mass was measured by ashing (500 °C for 2 h) the dried litter, cooling in a dessicator, re-weighing, and subtracting the ash weight (n=10).

Activities of three microbial extracellular enzymes (leucine aminopeptidase,  $\beta$ -glucosidase, and phosphatase) were compared at two levels of salinity: 2009.2  $\mu\text{S cm}^{-1}$  (ambient stream water) and  $\sim 60,000 \mu\text{S cm}^{-1}$ . Six replicates of the *Typha* litter sections and two litter-free blanks were used for each combination of salt treatment and enzyme type. Replicate litter sections (1.7 cm long) were pre-incubated in ambient or 33,040  $\mu\text{S cm}^{-1}$  stream water for 1 h (5°C, in darkness) before enzyme activity assays. The assays followed the method and conditions of Francoeur et al. (2006), which are saturating for these microbial communities. The containers containing the litter sections were aspirated of any liquid and 2 mL of 1000 uM stock solution, and one milliliter of 179,300  $\mu\text{S cm}^{-1}$  or 2009.2  $\mu\text{S cm}^{-1}$  stream water were added to each container. The final concentration of extracellular enzyme substrate was 666 uM stock solution, and final salt concentrations were approximately 60,000  $\mu\text{S cm}^{-1}$  (elevated salt treatment) or 2009.2  $\mu\text{S cm}^{-1}$  (ambient stream water treatment). Samples were incubated for 30 minutes in the dark at 5°C.

Aliquots from each vial were added to a 96-well plate with equal volumes of pH 10 buffer and measured fluorometrically .

*Experiment Three (March 13, 2007)*

On March 13, 2007, natural rock substrata were collected from the Paint Creek study site and returned to the lab with care to avoid losses of periphyton during transport. In the lab, replicate sections (~100 mm<sup>2</sup>) of intact periphyton mat were removed from the rock surfaces. Algal biomass (as chlorophyll a, n = 3) and community composition (n = 4) were quantified as in Experiment One. An additional 3 sections were preserved in 4% formalin and stored at 4°C in darkness for subsequent analysis of bacterial abundance. Bacterial samples were disrupted by probe sonication (Francoeur et al., 2006), stained with SYBR Gold, mounted on 0.2 µm Anodisc filters in anti-fade reagent (see Noble & Fuhrman, 1998), and examined using epifluorescence microscopy (1000×). At least 350 cells and 10 fields were examined for each sample.

Three salinity treatments were made by adding NaCl to stream water: 1609.8 µS cm<sup>-1</sup> (unamended stream water), 4428 µS cm<sup>-1</sup>, and 32750 µS cm<sup>-1</sup>. Seven replicate periphyton sections were immersed in 5.0 mL of each of the desired salt treatments. Incubation and photosystem II yield measurements were conducted as described above, with the exception that after 6 hours of incubation, all sections were removed from incubation water, rinsed with unamended stream water, and placed into unamended stream water. Incubation resumed and photosystem II yield measurements were made at 0.5 hours after rinsing, 3 hours after rinsing, and 21 hours from rinsing.

### *Statistical Analysis*

Experiment One. A Repeated-Measures ANOVA (RMANOVA) was conducted to test the hypothesis that potential algal photosynthetic activity (dark-adapted photosystem II yield) differed between ambient and elevated salt treatments over the 24 hour experiment. Additionally, two-sample separate-variance t-tests were used to test for differences between ambient and elevated salt treatments at each individual measurement period.

Experiment Two. Two-sample t-tests were used to test the hypothesis that activities of individual extracellular enzymes (leucine aminopeptidase,  $\beta$ -glucosidase, and phosphatase) differed between ambient and elevated salt treatments.

Experiment Three. RMANOVA was used to test for differences in dark-adapted photosystem II yield between salt treatments over the entire experiment. Bonferroni-based comparisons were used to identify differences between specific elevated salt treatments and the control. One-way ANOVAs with Tukey HSD multiple comparisons were used to test for differences among salt treatments at each individual measurement period. In addition, subsequent RMANOVAs considered only the pre-rinse measurements (0-6 h, when algae were exposed to three distinct salt treatments), and only the post-rinse measurements (6.5- 21.5 h, when all algae were in ambient salinity water). Finally, paired t-tests were used to test the hypothesis that dark-adapted photosystem II yield within individual salt treatments differed between the 6 h and 6.5 h measurements (i.e., immediately pre- and post-rinse).

Data were transformed as necessary to reduce heteroscedasticity. All statistical analyses were conducted using SYSTAT, Version 10 (Systat Software, Point Richmond, California).

## RESULTS

### *Environmental Conditions*

Over the course of the in situ incubations (January 4, 2007; January 18, 2007; January 29, 2007; February 2, 2007; February 22, 2007; March 6, 2007; March 13, 2007) the mean nutrient levels ( $\pm 1$  SE) in Paint Creek were  $2.29 \pm 0.56$  Nitrate-Nitrite mg/L ( $n=6$ );  $0.15673 \pm 0$   $\text{NH}_4^+$  mg/L ( $n=6$ );  $58.3 \pm 29.3$  TP  $\mu\text{g} /\text{L}$  ( $n=6$ ). Average stream water temperature was reported, and mean specific conductance was collected in situ or in the lab (Table 1). Algal biomass was greatest in the Experiment Three communities, moderate in the Experiment One communities, and lowest in the communities used in Experiment Two (Table 2). The only algae found were Bacilliarophyta. There were limited algal communities found in Experiment Two. The ash free dry mass mean ( $\pm$  SE) found within a sample taken on March 6 (Experiment Three) was  $0.05 (\pm 0.013)$   $\text{mg cm}^{-2}$ , and the dry mass mean ( $\pm$  SE) was  $0.03 (\pm 0.01)$   $\text{mg cm}^{-2}$ .

### *Photosynthesis Experiments*

In Experiment One, exposure to salt amended stream water ( $35,320 \mu\text{S cm}^{-1}$ ) significantly reduced photosynthetic yield of photosystem II ( $F_v/F_m$ ) over the entire 24-hour experimental period ( $p < 0.001$ , Figure 2) and at each individual sampling time ( $p$  always  $< 0.001$ ).

In regard to the full photosynthetic yield data set of Experiment Three, repeated-measures ANOVA detected significant differences among the three salinity levels ( $p < 0.001$ ) and a strong time x salinity interaction ( $p < 0.001$ ) (Figure 3). Both the medium ( $4,428 \mu\text{S cm}^{-1}$ ,  $p < 0.004$ ) and high ( $32,750 \mu\text{S cm}^{-1}$ ,  $p < 0.001$ ) treatments differ from the ambient-salinity control; however, the time x salinity interaction was only

apparent when comparing the high salinity and ambient treatments ( $p < 0.001$ ); the ambient and medium salinity treatments displayed the same pattern over time ( $p = 0.201$ ). Potential photosynthetic yield differed significantly among salinity treatments in every pre-rinse (i.e., times 0, 3, and 6) sampling period (one-way ANOVA  $p$  always  $< 0.001$ ).

A repeated-measures ANOVA considering only the pre-rinse sampling periods (i.e., times 0, 3, and 6) detected significant differences among salinity treatments ( $p < 0.001$ ); both high ( $p < 0.001$ ) and medium ( $p = 0.006$ ) salinity treatments differed from the ambient treatment. The high salinity treatment was significantly different from the ambient ( $p$  always  $< 0.001$ ) and medium salinity treatments ( $p$  always  $< 0.001$ ); however, the ambient and medium salinity treatments did not differ at individual pre-rinse sampling periods ( $p$  always  $> 0.424$ ).

Comparison of immediately pre-rinse (6 h) and immediately post-rinse (6.5 h) measurements showed that photosystem II yield in the ambient water ( $1609.8 \mu\text{S cm}^{-1}$ ) and low salinity ( $4,428 \mu\text{S cm}^{-1}$ ) treatments increased slightly, but the differences were not significant ( $p=0.071$  and  $p=0.797$ , respectively). However, the photosynthetic yield of the high salinity treatment ( $32,750 \mu\text{S cm}^{-1}$ ) increased significantly during this timeframe ( $p<0.001$ , Figure 3)

A repeated-measures ANOVA considering only the post-rinse sampling period did not differ significantly among salinity treatments ( $p = 0.060$ ). Potential photosynthetic yield in each of the post-rinse sampling periods (i.e., times 6.5, 9.5, and 21.5) reflected this similarity among salt treatments (one-way ANOVA  $p$  always  $< 0.086$ ).

### *Extracellular Enzyme Activity Experiment*

Extracellular enzyme activities (leucine aminopeptidase, phosphatase and beta-glucosidase) were not affected by high ( $\sim 60,000 \mu\text{S cm}^{-1}$ ) salinity ( $p = 0.413$ ,  $p=0.902$ ,  $p=0.899$ , respectively; Figure 4).

## DISCUSSION

The purpose of this study was to determine the short-term effects of salt stress on lotic microbial function in order to evaluate possible ecological effects from road salt run-off. We approached this task by exposing natural periphyton communities to simulated salting events. Laboratory conditions were calibrated to local environmental data in order to provide realistic salt concentrations and temperatures.

Potential photosynthetic activity showed a rapid reduction of activity when exposed to high, but realistic, salt concentrations. When high salt exposure samples were rinsed with ambient salinity stream water, a rapid recovery of photosynthetic activity was observed. In a Canadian gravel-bed river, Silva and Davis (1999) reported stimulation of natural lotic periphytic productivity by slightly increased salinity (up to  $\sim 6,000 \mu\text{S cm}^{-1}$ ), then strong reductions in periphytic productivity by salt stress in the highest salinity treatment ( $\sim 10,500 \mu\text{S cm}^{-1}$ ). In contrast to Silva and Davis (1999), I did not observe any stimulation of photosynthetic activity at medium levels of salinity ( $4,428 \mu\text{S cm}^{-1}$ ); the data indicated that these medium salinity levels had either no effect or were slightly inhibitory. This could have been from the osmotic stress from exposure to the concentration of the salt. Despite this inconsistency between the studies, high salinity results from this study are broadly similar to those of Silva and Davis (1999), despite their high salinity treatment being well below my high salt concentration ( $\sim 29,900 - 35,320 \mu\text{S cm}^{-1}$ ).

Not only was a salt-induced rapid decline observed in potential photosynthetic ability, but also a rapid recovery when high concentrations of salt were rinsed and replaced with unamended stream water. This general pattern of rapid decline and

recovery appears superficially similar to the results of Lu and Zhang (2000), who observed instantaneous decline, then recovery of photosynthetic activity of the cyanobacterium *Spirulina plantensis* from salt exposure. However, the recovery from salt stress in their experiment was spontaneous, as it did not require removal of salt from the medium, perhaps the result of sodium being actively excluded from the cyanobacterial cells (Lu & Zhang, 2000), whereas in my experiment, recovery was due to removal of algae from the salt solution. Lu and Zhang (2000) also reported that after 4 hours, there was a second, light-mediated depression of photosystem II activity by salt. This phenomenon was something that would not have been able to be observed in my experiment, as the samples were incubated in darkness. However, spontaneous recovery of photosynthetic activity during my experiments was not observed; thus, there is no way that a secondary light-mediated inhibition could have occurred. It is likely that the large differences in algal community composition between our experiments (natural algal communities almost exclusively composed of diatoms) and the purely cyanobacterial cultures of Lu and Zhang (2000) caused this disparity in spontaneous photosystem II recovery. The fact that the cyanobacterial spontaneous photosystem II recovery observed by Lu and Zhang (2000) was later reversed by light exposure strongly suggests that this apparent spontaneous recovery was dependent on the use of alternative photosynthetic electron pathways unique to cyanobacteria (Lu & Zhang, 2000). Such pathways are lacking in plants and eukaryotic algae (see Campbell et al., 1998), and thus this phenomenon would not be expected to occur in algal communities dominated by eukaryotic diatoms.

A more critical finding of Lu and Zhang (2000) is the decoupling of photosystem II activity and photosynthetic oxygen evolution by salt exposure. Should this decoupling be a common phenomenon, then rapid measures of photosystem II activity would not be good proxies for traditional photosynthesis assays. However, the discrepancy observed by Lu and Zhang (2000) appears to entirely result from the apparent spontaneous recovery of photosystem II activity. As discussed above, this phenomenon is likely limited to cyanobacteria. Furthermore, this apparent spontaneous recovery can be easily distinguished by its sensitivity to light. Exposure to  $100 \mu\text{mol m}^{-2} \text{s}^{-1}$  of light for several hours completely eliminated apparent spontaneous photosystem II recovery (Lu & Zhang, 2000). Thus, rapid, PAM-based measurement of photosystem II activity should generally provide an adequate estimate of eukaryotic algal photosynthetic performance under salt stress.

A similar decrease in photosynthetic oxygen production by *Synechococcus* sp. exposed to 0.5M NaCl was observed by Allakhverdiev et al. (2000). The cyanobacteria were rinsed after 0.5, 1 and 5 hours exposure to salt and the oxygen activity was measured. The samples that were rinsed after 0.5 hours of salt exposure were able to recover following a short exposure time to salt. The sample rinsed after 5 hours of salt exposure did not recover in their oxygen evolving activity of photosystem II. This lack of recovery is consistent with what Lu and Zhang (2000) found. In contrast, my study showed full recovery of photosynthetic activity in a diatom-dominated periphyton community following a six hour exposure of  $33,040 \mu\text{S cm}^{-1}$  salt. Taken together, these studies suggest that more work is required to understand the effects of salt on algae, as different species may react differently when exposed to various levels of salt.

In contrast to the strong responses observed for photosynthesis, extracellular enzyme activities showed no significant response to salt exposure. This could imply that there was no immediate effect of salt on decomposition. Enzymes are limited in what conditions they may function optimally within an environment and are limited by temperature, pH, concentration of ions, and salt (Audesirk et al., 2010; Tobin & Morel, 1997). Perhaps the biofilm nature of the extracellular enzymes buffered their activity against short-term salt exposure.

For determining the larger ecological implication of higher conductivity levels, a systems approach is necessary. Within Paint Creek, there were no rapid changes in the extracellular enzyme activity. However, potential photosynthetic activity was rapidly inhibited by high salt exposure but was also rapidly reversed by the removal of salt stress. This would suggest that if the salt exposure were temporary and short, there would be little to no long-term functional harm to stream microbes. Blasius & Merritt (2002) also support this finding using macroinvertebrates. Several species of macroinvertebrates were tolerant of short-term salt exposure. Furthermore, Silva & Davies (1999) also observed rapid recovery of macroinvertebrates following short-term exposure to salt. Rapid recovery of microbes following salt exposure was also observed in my study. However, there could be an effect on the production of other stream microbiota (e.g., bacteria, fungi, or protists), but we did not measure these other ecosystem components.

This study was designed to measure short-term physiological responses to salt exposure. Long-term salt exposure would be predicted to change species composition within periphyton communities. If the community were changed to include more halotolerant species because of the long-term exposure to salt, we could then expect some

change in ecological performance. Because of the very different mechanisms causing short-term (rapid physiological responses of existing taxa) and long-term (community shifts to halotolerant taxa) responses to salt exposure, it is difficult to extrapolate the results to long-term whole-stream production and decomposition. Little is known about the influence of long-term exposure of salt upon streams and benthic community development.

Lack of short-term effects on lotic microbial community function does not imply that short-term salt inputs are harmless to all freshwater ecosystems. In contrast to stream communities, where pulses of salt-rich water would be an acute problem during snowmelt, salt exposure in lake systems is a chronic impairment and can change physical, chemical, and biological processes of a lake. Increased salt concentrations introduced from road salt can create a lake with conductivity stratifications, thus preventing turnover from occurring in the spring and fall (Novotny et al., 2008). High density salt water accrues in the hypolimnion layer of the lake (Judd et al., 2005). Prevention of turnover has detrimental effects on a lake's water quality (Novotny et al., 2008). The turnover process in a lake helps the lake keep the benthic regions oxygenated, prevents anoxic regions from developing, and promotes natural nutrient availability (Judd et al., 2005; Novotny et al., 2008). An increased anoxic region within a lake in the fall reduces the extent of suitable habitat for fish in the winter (Novotny et al., 2008). Furthermore, altered nutrient availability may affect the community composition of the lake (Judd et al., 2005). If salt concentrations in the sediments of a lake increase, heavy metals may be released from the sediment into the water (Novotny et al., 2008). As chloride concentrations have correlated with increased salt concentrations in the state of

Minnesota since the 1950s, there appears to be a trend of decreasing health of lakes due to the application of road salt (Novotny et al., 2008).

Large but realistic short-term salt exposures immediately caused reductions in potential photosynthesis of lotic periphyton. Extracellular enzyme activities were not affected, and photosynthetic impairment was rapidly and completely reversible by rinsing with ambient, low-salt streamwater. Thus, individual short-term salt exposures do not appear to cause permanent harm to lotic periphyton.

## REFERENCES

- Allakhverdiev SI, Sakamoto A, Nishiyama Y, Inaba M, Murata N. 2000. Ionic and osmotic effects of NaCl-induced inactivation of Photosystems I and II in *Synechococcus* sp. *Plant Physiology* 123:1047-1056
- Allan JD, Castillo MM. 2007. *Stream Ecology: Structure and function of running waters*. 2<sup>nd</sup> edition. Springer, Dordrecht.
- Audesirk T, Audesirk G, Byers B. 2010. *Biology: Life on Earth*, 9th Edition. San Francisco, CA: Benjamin-Cummings.
- Biggs BJF, Kilroy C. 2000. *Stream Periphyton Monitoring Manual*. National Institute of Water and Atmospheric Research (NIWA), Christchurch, New Zealand.
- Blasius BJ, Merritt RW. 2002. Field and laboratory investigations on the effects of road-salt (NaCl) on stream macroinvertebrate communities. *Environmental Pollution* 120:219–231
- Blinn DW. 1993. Diatom community structure along physicochemical gradients in saline lakes. *Ecology* 74:1246–1263.
- Borchardt MA. 1996. Nutrients. RJ Stevenson, ML Bothwell, and RL Lowe (editors). *Algal Ecology: Freshwater Benthic Ecosystems*. Academic Press, San Diego (CA).
- Calow P. 1991. Physiological costs of combating chemical toxicants: Ecological implications. *Comparative Biochemistry and Physiology* 100:3–6.
- Campbell D, Hurry V, Clarke AK, Gustafsson P, Oquist G. 1998. Chlorophyll fluorescence analysis of cyanobacterial photosynthesis and acclimation. *Microbiology and Molecular Biology Reviews*. 62:667-683.
- Carr JM, Hergenrader GL, Troelstrup NH. 1986: A simple, inexpensive method for cleaning diatoms. *Transactions of the American Microscopical Society* 105:152-157.
- Clavero E, Hernandez-Marine M, Grimalt JO, Garcia-Pichel F. 2000. Salinity tolerance of diatoms from thalassic hypersaline environments. *Journal of Phycology* 36:1021-1034.
- Consalvey M, Perkins RG, Paterson DM, Underwood GJC. 2005. PAM fluorescence: A beginners guide for benthic diatoms. *Diatom Research* 20:1-22.
- Dodds WK, Whiles MR. 2010. *Freshwater Ecology: Concepts and environmental applications of limnology*. Academic Press, San Diego.

- Denoël M, Bichot M, Ficetola GF, Delcourt J, Yliff M, Kestemont P, Poncin P. 2010. Cumulative effect of road de-icing salt on amphibian behavior. *Aquatic Toxicology* 99:275–280.
- Dudley RJ, Churchill PF. 1995. Effect and potential ecological significance of the interaction of humic acids with two aquatic extracellular proteases. *Freshwater Biology* 34:485-494.
- Environment Canada. 2001. Canadian Environmental Protection Act, 1999: Priority Substance List Assessment Report Road Salts. Environment Canada, Hull, Quebec
- Francoeur SN, Schaecher M, Neely RK, Kuehn KA. 2006. Periphytic photosynthetic stimulation of extracellular enzyme activity in aquatic microbial communities associated with decaying *Typha* litter. *Microbial Ecology* 52:662-669.
- Francoeur SN, Wetzel RG. 2003. Regulation of periphytic leucine-aminopeptidase activity. *Aquatic Microbial Ecology* 31:249–258.
- Hauer R, Lamberti G. 1996. *Methods in Stream Ecology*. Academic Press. San Diego.
- Herbst DB, Blinn DW. 1998. Experimental mesocosm studies of salinity effects on the benthic algal community of a saline lake. *Journal of Phycology*. 34:772–778.
- Judd KE, Adams HE, Bosch NS, Kostrzewski JM, Scott CE, Schultz BM, Wang DH & Kling GW. 2005. A case history: Effects of mixing regime on nutrient dynamics and community structure in Third Sister Lake, Michigan during late winter and early spring 2003. *Lake and Reservoir Management* 21:316-329.
- Kahn WE, Wetzel RG. 1999. Effects of microscale water level fluctuations and altered ultraviolet radiation on periphytic microbiota. *Microbial Ecology* 38:253-263.
- Kelly VR, Lovett GM, Weathers KC, Findley SEG, Strayer DL, Burns DJ, Likens GE. 2008. Long-Term sodium chloride retention in a rural watershed: Legacy effects of road salt on stream water concentration. *Environmental Science and Technology* 42:410-415.
- Lind OT. 1985. *Handbook of common methods in limnology*, 2nd edition, Kendall/hunt, Dubuque, IA
- Lu C, Zhang J. 2000. Role of light in the response of PSII photochemistry to salt stress in the cyanobacterium *Spirulina platensis*. *Journal of Experimental Biology* 51:911-917.
- Noble RT, Fuhrman JA. 1998. Use of SYBR Green I for rapid epifluorescence counts of

- marine viruses and bacteria. *Aquatic Microbial Ecology* 14:113-118.
- Novotny EV, Murphy D, Stefan HG. 2008. Increase of urban lake salinity by road deicing salt. *Science of the Total Environment* 406:131–144
- Oskinis V, Kasperovicus T. 2005. Impact of road maintenance salts on water ecosystems according to diatom flora investigation. *Journal of Environmental Engineering and Landscape Management* 13:51-55.
- Paschka MG, Ghosh RS, Dzombak DA. 1999. Potential water-quality effects from iron cyanide anticaking agents in road salt. *Water Environment Research* 71:1235-1239.
- Pohlon E, Marxsen J, Küsel K. 2010. Pioneering bacterial and algal communities and potential extracellular enzyme activities of stream biofilms. *FEMS Microbiology Ecology* 71:364-373.
- Ramakrishna DM, Viraraghavan T. 2005. Environmental impact of chemical deicers - a review. *Water, Air, and Soil Pollution* 166:49-63.
- Rier ST, Kuehn KA, Francoeur SN 2007. Algal regulation of extracellular enzyme activity in stream microbial communities associated with inert substrata and detritus. *Journal of the North American Benthological Society* 26:439-449.
- Rier ST, Nowrocki KS, Whitley JC. 2011. Response of biofilm extracellular enzymes along a stream nutrient enrichment gradient in an agricultural region of north central Pennsylvania, USA. *Hydrobiologia* 669:119-131.
- Sanzo D, Hecnar SJ. 2006. Effects of road de-icing salt (NaCl) on larval wood frogs (*Rana sylvatica*). *Environmental Pollution* 140:247–256.
- Saros JE, Fritz SC. 2000. Nutrients as a link between ionic concentrations/composition and diatom distributions in saline lakes. *Journal of Paleolimnology* 23:449– 453.
- Silva EIL, Davies RW. 1999 The effects of simulated irrigation induced changes in salinity on metabolism of lotic biota. *Hydrobiologia* 416:193-202.
- Silva EIL, Shimizu A, Matsunami H. 2000. Salt pollution in a Japanese stream and its effects on water chemistry and epilithic algal chlorophyll-*a*. *Hydrobiologia*. 437:139-148.
- Sinsabaugh RL, Linkins AE. 1993. Statistical modeling of litter decomposition from integrated cellulase activity. *Ecology* 74:1594–1597.
- Sinsabaugh RL, Osgood MP, Findlay S. 1994. Enzymatic models for estimating decomposition rates of particulate detritus. *Journal of the North American*

- Benthological Society 13:160–169.
- Sinsabaugh RL, Van Horn DJ, Follstad Shah JJ, Findlay S. 2010. Ecoenzymatic stoichiometry in relation to productivity for freshwater biofilm and plankton communities. *Microbial Ecology* 60:885-893.
- Sudhir P, Pogorylelov D, Kovacs L, Garab G, Murthy SDS. 2005. The effects of salt stress on photosynthetic electron transport and tylakoid membrane proteins in the cyanobacterium *Spirulina platensis*. *Journal of Biochemistry and Molecular Biology*. 38:481-485.
- Tobin AJ, Morel RE 1997. *Asking About Cells*, Saunders College Publishing, USA.
- Washtenaw County Road Commission Website. [www.wcroads.org](http://www.wcroads.org). Last Accessed on 11-07-06.
- Williams DD, Williams NE., Cao Y. 1999. Road salt contamination of groundwater in a major metropolitan area and development of a biological index to monitor its impact. *Water Research* 34:127-138.


**Table 1.** Environmental conditions at the Paint Creek field site. Values are means ( $\pm 1$  SD) of six measurements made from 4 January – 13 March 2007). Asterisk indicates water samples were taken from stream and water temperature and conductance were measured upon returning to lab. n.d. = not determined.

	January 4, 2007	January 18, 2007	January 29, 2007	February 2, 2007	February 6, 2007 (Experiment 1)	February 22, 2007	March 6, 2007 (Experiment 2)	March 13, 2007 (Experiment 3)
<b>Water Temperature (°C)</b>	4.82 $\pm$ 0.04	3.7	0.65 $\pm$ 0.07	1.76 $\pm$ 0.37	3.6 $\pm$ 0.12	3.6 $\pm$ 0.12	5.68 $\pm$ 0.16**	12.46 $\pm$ 0.05**
<b>Specific Conductance (<math>\mu</math>S/cm)</b>	852.8 $\pm$ 1.9	1377.4 $\pm$ 20.0	n.d.	1805.8 $\pm$ 10.3	1637.2 $\pm$ 13.4	2256.6 $\pm$ 11.3	2009.2 $\pm$ 2.5**	1609.8 $\pm$ 3.2**
<b>NO<sub>3</sub>+NO<sub>2</sub> (mg/L)</b>	2.67	2.99	2.30	1.49	n.d.	1.43	n.d.	1.83
<b>NH<sub>4</sub> (mg/L)</b>	0.159	0.160	0.156	0.140	n.d.	0.165	n.d.	0.160
<b>TP (<math>\mu</math>g/L)</b>	105	65	25	30	n.d.	55	n.d.	70


**Table 2.** Mean ( $\pm 1$  SE) algal biomass, fungal biomass, and bacterial abundance of periphyton communities used for the three experiments. n.d. = not determined.

	<b>February 6, 2007</b>	<b>March 6, 2007</b>	<b>March 13, 2007</b>
<b>Algal Biomass (mg Chl a / m<sup>2</sup>) (n=3)</b>	37 $\pm$ 6	16 $\pm$ 8	86 $\pm$ 12
<b>Fungal Biomass (mg/g dwt.) (n = 4)</b>	n.d.	53.15 $\pm$ 7.6	n.d.
<b>Bacterial Abundance (x 10<sup>10</sup> cells L<sup>-1</sup>) (n = 3)</b>	n.d.	n.d.	5.89 $\pm$ 4.47


**Figure 1.** Conductivity measurements ( $\mu\text{S cm}^{-1}$ ,  $n = 1456$ , 1995-2006) from Huron River tributaries in Washtenaw County, MI and surrounding region. Data are sorted highest to lowest, and are courtesy of the Huron River Watershed Council (HRWC.org).



**Figure 2.** Effects of conductivity on dark-adapted photosystem II yield (Fv/Fm) in experiment one ( $p < 0.001$ ). Values are means  $\pm$  1 SE (n=7).



**Figure 3.** Effects of conductivity on dark-adapted photosystem II yield (Fv/Fm) in experiment three. Photosystem II yield measurements means were significantly different before a rinse ( $p < 0.001$ ), but not significantly different following the rinse at 6 hours and before 6.5 hours ( $p = 0.061$ ).



**Figure 4.** Mean extracellular enzyme activity ( $\text{nmol cm}^{-2} \text{h}^{-1}$ ) in conductivity treatments at ambient stream water ( $2009.2 \mu\text{S cm}^{-1}$ ) and salt-amended stream water ( $\sim 60,000 \mu\text{S cm}^{-1}$ ).

